



REPLY TO LETTER

Reply to: "Inappropriate interpretation of non-pathogenic *HTRA1* variant as pathogenic"

Chen Zhang^{1,2} , Honghua Zheng³, Xin Li³, Shaowu Li⁴, Wei Li^{2,5}, Ziwei Wang³, Songtao Niu^{1,2}, Xingao Wang^{1,2} & Zaiqiang Zhang^{1,2} 

¹Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

²National Clinical Research Center for Neurological Diseases, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

³Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, Institute of Neuroscience, School of Medicine, Xiamen University, Fujian, Xiamen, China

⁴Beijing Neurosurgical Institute, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

⁵Monogenic Disease Research Center for Neurological Disorders, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

Dear Editor,

We thank Uemura et al.¹ for their letter suggesting interpreting the *HTRA1* A20V variant as non-pathogenic.

It is true that the *HTRA1* A20V variant had a higher frequency of carriers compared with the other pathogenic *HTRA1* variants identified in our study. However, it must be noted that all carriers with *HTRA1* A20V variants in our study had clinical phenotypes that were consistent with *HTRA1*-related cerebral small vessel disease. Moreover, carriers with both homozygous and heterozygous *HTRA1* A20V variants were affected, and family members without this variant were not affected.²

Many of the pathogenic *HTRA1* variants reported to date are positioned within domains known to affect enzymatic function. The p.A20V (c.59C>T) variant is positioned within *HTRA1*'s N-terminal domain, the function(s) of which remain largely unknown.³ It does bear mention that a study reported autolysis of the N-terminal domain does not affect protease activity.⁴ Functional investigations into the pathophysiological impacts of the p.A20V (c.59C>T) variant will be necessary to better understand its role in the development of cerebral small vessel disease.

Funding Information

This study was supported by a grant from the Beijing Municipal Science and Technology Commission (No. Z171100001017080).

Conflict of Interest

None.

References

1. Uemura M, Kitahara S, Nozaki H, et al. Inappropriate interpretation of non-pathogenic *HTRA1* variant as pathogenic. *Ann Clin Transl Neurol*. 2023;In press.
2. Zhang C, Zheng H, Li X, et al. Novel mutations in *HTRA1*-related cerebral small vessel disease and comparison with CADASIL. *Ann Clin Transl Neurol*. 2022;9:1586-1595.
3. Truebestein L, Tennstaedt A, Monig T, et al. Substrate-induced remodeling of the active site regulates human *HTRA1* activity. *Nat Struct Mol Biol*. 2011;18:386-388.
4. Risor MW, Poulsen ET, Thomsen LR, et al. The autolysis of human *HtrA1* is governed by the redox state of its N-terminal domain. *Biochemistry*. 2014;53:3851-3857.